

## 3-Year Canaloplasty & Combined<sup>1</sup>:

### At 36 Months

#### Canaloplasty Only

- Mean decrease in IOP from Baseline **34%**
- Mean reduction in Medication **53%**

#### PhacoCanaloplasty

- Mean decrease in IOP from Baseline **42%**
- Mean reduction in Medication **80%**

Data Point	Baseline	6 Months	12 Months	24 Months	36 Months
<b>Group 1: Canaloplasty Only</b>					
N	103	86	91	89	89
Mean IOP	<b>23.5 ± 4.5</b>	16.1 ± 3.8	16.1 ± 3.9	16.1 ± 4.0	<b>15.5 ± 3.5</b>
Mean Medications	<b>1.9 ± 0.8</b>	0.4 ± 0.7	0.6 ± 0.8	0.6 ± 0.8	<b>0.9 ± 0.9</b>
<b>Group 2: PhacoCanaloplasty</b>					
N	30	25	27	25	27
Mean IOP	<b>23.5 ± 5.2</b>	12.8 ± 2.9	13.6 ± 4.1	13.4 ± 3.2	<b>13.6 ± 3.6</b>
Mean Medications	<b>1.5 ± 1.0</b>	0.1 ± 0.3	0.1 ± 0.4	0.2 ± 0.4	<b>0.3 ± 0.5</b>

References: 1. Lewis RA, von Wolff K, Tetz M, et al. Canaloplasty: Three-year results of circumferential viscodilation and tensioning of Schlemm's canal using a microcatheter to treat open-angle glaucoma. J Cataract Refract Surg 2011 [in press].